

## IBD sharing around the *PPARG* locus is not increased in dizygotic twins or their mothers

Busjahn *et al*<sup>1</sup> reported finding linkage between *PPARG* (3p25) and being a dizygotic (DZ) twin. We differentiate, as do the authors, between the conception of DZ twins, and being the viable result of such a conception.

We observe no evidence of linkage to the region around the *PPARG* locus in

several samples of DZ twins who have been genotyped at multiple markers on chromosome 3 (Fig. 1). Among 199 Australian DZ twins ascertained for a history of wheezing<sup>2</sup>, mean identity by descent (IBD) sharing at the position of *PPARG* is 0.463 (99% bootstrapped confidence interval=0.412–0.516). We obtained a

similar result with 232 pairs of Australian adolescent DZ twins taking part in a longitudinal study of naevus development<sup>3</sup> (0.444, 0.390–0.499), and a set of 125 Australian adult DZ twin pairs assessed for anxiety<sup>4</sup> (0.508, 0.435–0.580). A Dutch scan of 160 DZ twin pairs<sup>5</sup> obtained slightly more encouraging results (0.553, 0.482–0.587, peak maximum lod score (MLS)=0.57). Pooling all these samples gives 0.477 (0.454–0.512) at the position of *PPARG*. The test for heterogeneity of sharing between studies was not significant ( $P=0.10$ ). In the combined dataset, the peak IBD sharing (MLS=0.70) is 50 cM closer to the centromere than *PPARG*.

Finally, in a sample of 203 Australian and New Zealand sister pairs where each had given birth to DZ twins<sup>6</sup>, sharing across the region is also not increased (0.433). We do not replicate linkage in the populations we study to survival of a twin pregnancy or polyovulation.

David Duffy<sup>1</sup>, Grant Montgomery<sup>1</sup>, Susan Treloar<sup>1</sup>, Andrew Birley<sup>1</sup>, Katherine Kirk<sup>1</sup>, Dorret Boomsma<sup>2</sup>, Leo Beem<sup>2</sup>, Eco de Geus<sup>2</sup>, Eline Slagboom<sup>3</sup>, Johanna Knighton<sup>4</sup>, Peter Reed<sup>4</sup> & Nicholas Martin<sup>1</sup>

<sup>1</sup>The Queensland Institute of Medical Research, Brisbane, Australia. <sup>2</sup>Free University, Amsterdam, The Netherlands. <sup>3</sup>Gaubius Laboratory TNO/PG, Leiden, The Netherlands. <sup>4</sup>Gemini Genomics, Cambridge, UK.

Received 4 January; accepted 27 June 2001.

1. Busjahn, A. *et al.* *Nature Genet.* **26**, 398–399 (2000).
2. Duffy, D. *et al.* *Am. Rev. Respir. Crit. Care Med.* **157**, 840–845 (1998).
3. Zhu, G. *et al.* *Am. J. Hum. Genet.* **65**, 483–492 (1999).
4. Boomsma, D. *et al.* *Twin Res.* **3**, 323–334 (2000).
5. Kirk, K. *et al.* *Twin Res.* **3**, 299–309 (2000).
6. Montgomery, G. *et al.* *J. Clin. Endocrinol. Metab.* **85**, 3391–3395 (2000).




Fig. 1 Mean IBD sharing from five studies of DZ twins and twinning.